

## ASSIGNMENT #5

**Please do not write your answers on a copy of this assignment.** As with all assignments, there will be conceptual and computational questions. For computational problems you may check your work using any tool you wish; however **you must clearly explain each step that you make in your computation.**

For this assignment I encourage you to work with others; however, you are expected to **submit your own work in your own words.** In addition to the true and false section being graded, I will grade one other problem; this will account for 10 points out of 25. The other 15 will be based on completion. **If you would like feedback on a particular problem, please indicate it somehow.** You must make an honest attempt on each problem for full points on the completion aspect of your grade.

(1) Answer each of the following questions. Also, the following parts are not related to one another.

(a) Let  $A = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix}$  and define  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  by  $T(\mathbf{x}) = A\mathbf{x}$ . Find the images of  $\mathbf{u} = \begin{bmatrix} 1 \\ -3 \end{bmatrix}$  and  $\mathbf{v} = \begin{bmatrix} a \\ b \end{bmatrix}$  under  $T$ .

(b) Let  $A = \begin{bmatrix} 1 & -2 & 1 \\ 3 & -4 & 5 \\ 0 & 1 & 1 \\ -3 & 5 & 4 \end{bmatrix}$  and  $\mathbf{b} = \begin{bmatrix} 1 \\ 9 \\ 3 \\ -6 \end{bmatrix}$ . Define  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^4$  by  $T(\mathbf{x}) = A\mathbf{x}$ . Find a vector  $\mathbf{x}$  such that  $T(\mathbf{x}) = \mathbf{b}$ .

(c) Let  $A = \begin{bmatrix} 1 & 2 & 2 & 1 \\ 7 & 1 & 1 & 2 \end{bmatrix}$ . Define  $T : \mathbb{R}^4 \rightarrow \mathbb{R}^2$  by  $T(\mathbf{x}) = A\mathbf{x}$ . Find all vectors  $\mathbf{x} \in \mathbb{R}^4$  such that  $T(\mathbf{x}) = \mathbf{0}$ .

(d) Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a linear transformation that maps  $\mathbf{u} = \begin{bmatrix} 5 \\ 2 \end{bmatrix}$  to  $\begin{bmatrix} 2 \\ 1 \end{bmatrix}$  and  $\mathbf{v} = \begin{bmatrix} 1 \\ 3 \end{bmatrix}$  to  $\begin{bmatrix} 1 \\ -3 \end{bmatrix}$ . Using what it means for  $T$  to be a linear transformation, find  $T(3\mathbf{u})$ ,  $T(2\mathbf{v})$ , and  $T(3\mathbf{u} + 2\mathbf{v})$ .

(2) Assume  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  is a linear transformation. For each situation below find a matrix that represents  $T$ .

(a)  $T$  rotates points about the origin by  $-\frac{\pi}{4}$  radians

(b)  $T$  is a vertical shear that maps  $\mathbf{e}_1$  to  $3\mathbf{e}_1 + 2\mathbf{e}_2$  but leaves  $\mathbf{e}_2$  unchanged (i.e.  $T(\mathbf{e}_2) = \mathbf{e}_2$ ). Recall that  $\mathbf{e}_1 = (1, 0)$  and  $\mathbf{e}_2 = (0, 1)$

(c)  $T$  first reflects points through the vertical  $x_2$  axis and then rotates points by  $\frac{3}{2\pi}$  radians about the origin. *Hint:* You might consider each transformation one at a time, and then multiply the matrices you find.

(3) For each of the matrices below with \*'s for entries, fill in the \*'s with numbers that make each a true statement.

$$(a) \begin{bmatrix} * & * & * \\ * & * & * \\ * & * & * \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 2x_1 - 3x_3 \\ 4x_1 \\ x_1 - x_2 + x_3 \end{bmatrix}.$$

$$(b) \begin{bmatrix} * & * \\ * & * \\ * & * \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} x_1 - 3x_2 \\ x_1 \\ 7x_2 \end{bmatrix}.$$

(c) Find a matrix that realizes the linear transformation  $T : \mathbb{R}^4 \rightarrow \mathbb{R}^2$ , where

$$T(x_1, x_2, x_3, x_4) = (2x_1 + x_3, x_2 - x_4).$$

*Hint:* What you did for parts (a) and (b) may be helpful here.

$$(4) \text{ Let } A_1 = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}, A_2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}, \text{ and } \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}.$$

(a) Find all vectors in  $\mathbb{R}^3$  that are fixed by  $A_1$ . In other words, find all  $\mathbf{x} \in \mathbb{R}^3$  such that  $A_1\mathbf{x} = \mathbf{x}$

(b) Find all vectors in  $\mathbb{R}^3$  that are fixed by  $A_2$ . In other words, find all  $\mathbf{x} \in \mathbb{R}^3$  such that  $A_2\mathbf{x} = \mathbf{x}$

(c) Find all vectors in  $\mathbb{R}^3$  that are fixed by  $A_3$ . In other words, find all  $\mathbf{x} \in \mathbb{R}^3$  such that  $A_3\mathbf{x} = \mathbf{x}$

(d) Use parts (a) through (c) to find all vectors in  $\mathbb{R}^3$  that are fixed by  $A_1, A_2, A_3$ .

(e) Congratulate yourself! You have done a little bit of representation theory! These matrices generate the group of permutations of a three element set. This exercise is part of the steps you would take to find all irreducible representations of this permutation group (whatever that means).

(5) Suppose that  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  is a linear transformation such that  $T(\mathbf{e}_1) = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ ,  $T(\mathbf{e}_2) = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}$ , and

$T(\mathbf{e}_3) = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$ . Answer the following questions.

(a) Find a matrix  $A$  that represents  $T$ . In other words, find a matrix  $A$  such that  $T(\mathbf{x}) = A\mathbf{x}$  for all  $\mathbf{x} \in \mathbb{R}^3$ .

(b) Describe the set of vectors,  $\mathbf{x}$ , that satisfy  $A\mathbf{x} = \mathbf{0}$  in vector parametric notation.

(c) Does  $A$  have an inverse? If so, what is it?

(d) Is  $T$  an isomorphism? Why or why not?

- (6) Answer the following as true or false. No justification is needed.
- (a) Every matrix determines a linear transformation.
  - (b) Every linear transformation is determined by a matrix.
  - (c) There is a linear transformation  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$  that is surjective.
  - (d) There is a linear transformation  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$  that is injective.
  - (e) If  $T : \mathbb{R}^m \rightarrow \mathbb{R}^n$  and  $S : \mathbb{R}^p \rightarrow \mathbb{R}^m$  are linear transformations, then  $T \circ S : \mathbb{R}^p \rightarrow \mathbb{R}^n$  is a linear transformation.
  - (f) If  $T$  is a linear transformation given by a  $n \times m$  matrix, then its codomain is  $\mathbb{R}^n$ .